

Equine Vaccination Protocols

<p>**denotes vaccines to be considered "Core" meaning all animals are at risk and should be vaccinated, all other protocols listed are risk-based and should be discussed with your veterinarian</p>	<p>Foals and Weanlings (<12 months of age) of mares vaccinated in the prepartum period against the disease indicated</p>	<p>Foals and Weanlings (<12 months of age) of unvaccinated mare or lacking vaccination history</p>	<p>Broodmares</p>	<p>Other Adult Horses (>1 year of age) previously vaccinated against the disease indicated</p>	<p>Other Adult Horses (>1 year of age) unvaccinated or lacking vaccine history</p>
<p>Tetanus** <u>can be given by itself or part of Prestige V+WNV</u> *Booster at time of penetrating injury or prior to surgery if last dose was administered over 6 months previously.</p>	<p>3 dose series: 1st dose- 4-6 months of age 2nd dose- 4-6 weeks after the 1st dose 3rd- 10-12 months of age</p>	<p>3 dose series: 1st dose- 3-4 months of age 2nd dose- 4-6 weeks after the 1st dose 3rd dose- 10-12 months of age</p>	<p><i>Previously vaccinated:</i> Annual, 4-6 weeks prior to foaling <i>Previously unvaccinated or unknown history: 2 dose series- 2nd dose 4-6 weeks after the 1st dose</i></p>	<p>Annual</p>	<p>2 dose series: 2nd dose 4-6 weeks after the 1st dose, annual revaccination</p>
<p>Eastern/Western Equine Encephalomyelitis** (<i>Sleeping Sickness</i>) <u>Part of Prestige V+WNV</u> *Consider 6 month revaccination interval for immunocompromised horses</p>	<p>3 dose series: 1st dose: 4-6 months of age 2nd dose: 4-6 weeks after the 1st dose 3rd- 10-12 months of age</p>	<p>3 dose series: 1st dose- 3-4 months of age 2nd dose- 4 weeks after 1st dose 3rd dose- 8 weeks after 2nd dose</p>	<p><i>Previously vaccinated:</i> Annual, 4-6 weeks prior to foaling <i>Previously unvaccinated or unknown history: 2 dose series- 2nd dose 4 weeks after the 1st dose</i></p>	<p>Annual- best given in spring</p>	<p>2 dose series: 2nd dose 4-6 weeks after the 1st dose, annual revaccination- best if given in spring</p>
<p>Rabies**</p>	<p>1 dose- 6 months of age Annual Revaccination</p>	<p>2 dose series: 1st dose- 6 months of age 2nd dose-4-6 weeks after 1st dose</p>	<p>4-6 weeks prior to foaling OR prior to breeding due to relatively long duration of immunity</p>	<p>Annual</p>	<p>Single dose, annual revaccination</p>

<p>West Nile Virus** <u>Part of Prestige+WNV</u></p>	<p>3 dose series: 1st dose: 4-6 months of age 2nd dose: 4-6 weeks after the 1st dose 3rd- 10-12 months of age</p>	<p>3 dose series: 1st dose- 3-4 months of age 2nd dose- 4 weeks after 1st dose 3rd dose- 8 weeks after 2nd dose</p>	<p><i>Previously vaccinated:</i> Annual, 4-6 weeks prior to foaling <i>Previously unvaccinated or unknown history:</i> it is preferable to vaccinate naïve mares when open</p>	<p>Annual- best given in spring</p>	<p>2 dose series: 2nd dose 4-6 weeks after the 1st dose, annual revaccination- best if given in spring</p>
<p>Strangles <u>Pinaccele I.N.</u> *Vaccination is not recommended as a strategy in outbreak mitigation</p>	<p>3 dose series administered intranasally: 1st dose: 6-9 months of age 2nd dose: 3-4 weeks after the 1st dose 3rd- 11-12 months of age</p>	<p>3 dose series administered intranasally: 1st dose: 6-9 months of age 2nd dose: 3-4 weeks after the 1st dose 3rd- 11-12 months of age</p>	<p><i>Previously vaccinated:</i> Killed vaccine containing M protein: semi- annual, 4-6 weeks prior to foaling <i>Previously unvaccinated or unknown history:</i> Killed vaccine containing M protein: 3 dose series 2nd dose 2-4 weeks after 1st dose, 3rd dose 4-6 weeks prior to foaling</p>	<p>Semi-annual to annual</p>	<p>Modified live vaccine: 2-dose series administered intranasally: 2nd dose 3 weeks after 1st dose, revaccinate annually to semi-annually</p>
<p>Influenza <u>Part of the Prestige V +WNV</u></p>	<p>3 dose series: 1st dose: 6 months of age 2nd dose: 3-4 weeks after 1st dose 3rd dose: 10-12 months of age</p>	<p>3 dose series: 1st dose: 6 months of age 2nd dose: 3-4 weeks after 1st dose 3rd dose: 10-12 months of age</p>	<p><i>Previously vaccinated:</i> Semiannual, 4-6 weeks prior to foaling <i>Previously unvaccinated or unknown history:</i> 3 dose series: 2nd dose 4-6 weeks after 1st dose, 3rd dose 4-6 weeks prior to foaling</p>	<p>Semi-annual to annual</p>	<p>3 dose series: 2nd dose 4-6 weeks after 1st dose, 3rd dose 3-6 months after 2nd dose, revaccinate semi-annually to annually</p>
<p>Equine Herpes Virus <u>Pneumabort</u></p>			<p>3 dose series: give at 5, 7, and 9 months of gestation</p>		